

Diffuse X-Ray Scattering Study of the Local Atomic Order and Pair Displacements in a $\text{Au}_{50}\text{Cu}_{50}$

G. Ice, C. Sparks, J. Bai (ORNL) and O. Malis (ORNL/Cornell U.)

Abstract No. ice7835

Beamline(s): X15A

Earlier diffuse x-ray scattering measurements on Au-Cu_3 alloy¹ with a single x-ray energy concluded that Au-Au first neighbor pairs are on average closer together than Au-Cu or Cu-Cu pairs. This conclusion is contrary to intuition as the Au atom is ~13% larger than the FCC Cu lattice and cast doubt on the sensitivity of diffuse x-ray scattering to measure atom size. Also, the addition of Au to Cu expands the lattice and ordering of AuCu_3 (which makes more Au-Cu first neighbor pairs) decrease the lattice parameter. Theoretical considerations and EXAFS measurements support the notion of larger Au-Au and smaller Cu-Cu first neighbor bond distances. Because of our advancements in the use of energy tunable synchrotron radiation to highlight specific elements in diffuse x-ray scattering measurements and to separate the individual specific pair bond distances, we undertook these measurements to show their application to a wide variety of solid solutions with atomic numbered elements both close and widely separated.²⁻⁶

Three dimensional volume measurements with x-ray energies chosen to be 20 eV below the Cu K edge and Au L_3 edge and an energy in-between reveal the asymmetry in the short-range order maxima associated with the variations in the neighboring atom bond distances. Diffuse x-ray data for a plane in reciprocal space is shown in Figure 1. Analysis is underway to recover the individual pair displacements.

References

1. B. D. Butler and J. B. Cohen, *J. Appl. Phys.* **65**(6), 2214 (1989).
2. G. E. Ice, C. J. Sparks, A. Habenschuss, and L. B. Shaffer, *Phys. Rev. Lett.* **68**, 863 (1992).
3. L. Reinhard, J. L. Robertson, S. C. Moss, G. E. Ice, P. Zschack, and C. J. Sparks, *Phys. Rev. B* **45**, 2662 (1992).
4. B. Schönfeld, G. E. Ice, C. J. Sparks, H. G. Haubold, W. Schweika, and L. B. Shaffer, *Phys. Status Solidi B* **183**, 79 (1994).
5. G. E. Ice, C. J. Sparks, and L. Shaffer, in *Resonant Anomalous X-Ray Scattering: Theory and Experiment*, edited by G. Materlik, C. J. Sparks, and K. Fischer (North-Holland, Amsterdam, 1994), p. 265.
6. J. L. Robertson, G. E. Ice, C. J. Sparks, X. Jiang, P. Zschack, F. Bley, S. Lefebvre, and M. Bessiere, *Phys. Rev. Lett.* **82**(14), 2911 (1999).




Figure 1. A planer cut through reciprocal space at $L=0.2$ for a $\text{Au}_{50}\text{Cu}_{50}$ single crystal quenched from 475°C . At the x-ray energy of 8.964 keV, the maximum contrast between Au and Cu atoms is obtained. The short-range order maxima [near (2,1), (3,1), (4,1), (3,2), (3,3)] show a systematic displacement (asymmetry) associated with the different atom pair bond distances.